

**THE NUMBER OF MINIMAL LATTICE PATHS
RESTRICTED BY TWO PARALLEL LINES****Masako SATO***Department of Mathematical Sciences, College of Engineering, University of Osaka Prefecture,
Sakai, Osaka, 591, Japan***Thien Tran CONG***Nippon Computer System Corporation, Sumitomo Nakanoshima Bldg. 10F, 3-2-18, Nakanoshima
Kita-ku, Osaka 530, Japan*

Received 11 September, 1981

We deal with non-decreasing paths on the non-negative quadrant of the integral square lattice, called by minimal lattice paths, from $(0, 0)$ to a point (n, m) restricted by two parallel lines with an incline k (≥ 0). We express the generating functions of the number of these distinct minimal lattice paths in terms of the polynomials

$$\varphi_k(n, x) = \sum_{l=0}^{\lfloor n/k \rfloor} \binom{n-kl}{l} (-x)^l, \quad n \geq 0.$$

Formulas obtained thus include the generating function of the so-called higher Catalan number $C_k(n)$ or Ballot numbers as the special case.

The number of minimal lattice paths for $k=1$ is given as an explicit form by expanding the corresponding generating function.

1. Introduction

Consider a particle executing a random walk on the interval $[-b, a]$ where a and b are positive integers. The particle in a point j ($-b < j < a$) moves to one of the points $j-1$ and $j+1$ in a single transition. Both of the points a and $-b$ are taken to be absorbing barriers. First, we consider random walk paths in which the particle arrives at a point i ($-b < i < a$) after t transitions, starting from an initial point 0 without touching any of the absorbing barriers a and $-b$. If the paths consist of n -right moves and m -left moves, then each of these paths corresponds to a non-decreasing path on the nonnegative quadrant of the integral square lattice, called minimal lattice path hereafter, from $(0, 0)$ to the point (n, m) , without crossing any of the lines $y = x - a + 1$ and $y = x + b - 1$, where $n + m = t$ and $n - m = i$. The number of distinct paths satisfying these restrictions is given by (S. Dua et al. [1])

$$\sum_{l=-\infty}^{\infty} \left\{ \binom{t}{\frac{1}{2}(t+i)+l(a+b)} - \binom{t}{\frac{1}{2}(t+i)+b+l(a+b)} \right\}, \quad t+i \equiv 0 \pmod{2} \quad (1)$$

with the usual convention that $\binom{t}{n} = 0$ for $n > t$ or $n < 0$.

In this present paper, we deal with generating functions of the following numbers $\{W_k(N/k, n, r), n \geq 0\}$, and $\{T_k(N/k, n, r, s), n \geq 0\}$, respectively:

(i) $W_k(N/k, n, r)$ is the number of minimal lattice paths from $(0, 0)$ to $(n, kn - N + r)$ without crossing a line $y = k(x - N/k)$ (see Fig. 1),

(ii) $T_k(N/k, n, r, s)$ is the number of minimal lattice paths from $(0, 0)$ to $(n, kn - N + r)$ without crossing any of lines $y = k(x - N/k)$ and $y = kx + s$ (see Fig. 2), where $n, N, r, s \geq 0$ and $k \geq 1$.

According to the above notations, the finite sum (1) is equal to $T_1(a-1, n, a-1-i, b-1)$, where $n = \frac{1}{2}(t+i)$ and $t+i \equiv 0 \pmod{2}$.

The number $W_k(0, n, 0)$ of minimal lattice paths from $(0, 0)$ to (n, kn) without crossing the line $y = kx$ is the well known higher Catalan number (Rogers [2])

$$C_k(n) = \frac{1}{kn+1} \binom{(k+1)n}{n}, \quad n \geq 0.$$

The higher Catalan number $C_k(n)$, like the Catalan number $C_1(n)$, occurs in a wide variety of combinatorial problems [2-5]. The number $W_k(0, n, r)$ of minimal lattice paths from $(0, 0)$ to $(n, kn + r)$ without crossing the line $y = kx$ is also





Fig. 1. $W_k(N/k, n, r)$.

Fig. 2. $T_k(N/k, n, r, s)$.

represented by (Rogers [2])

$$W_k(0, n, r) = A_n(r+1, k+1) = \frac{r+1}{r+1+(k+1)n} \binom{r+1+(k+1)n}{n}, \quad n \geq 0$$

with $A_n(0, k+1) = C_k(n)$.

In our previous paper [6], we have expressed $W_k(N/k, n, r)$ in terms of $\{A_l(r+1, k+1)\}$ for $N, n, r \geq 0$ and $kn-N+r \geq 0$. In Section 2, we give the generating functions for $\{W_k(N/k, n, r), n \geq 0\}$. In Section 3, we express $T_k(N/k, n, r, s)$ in terms of $\{W_k(N'/k, l, r')\}$. After that, we obtain the generating functions for $\{T_k(N/k, n, r, s), n \geq 0\}$ where $N, s \geq 0$ and $0 \leq r \leq s+N$. It is proved that the generating function for $\{T_k(N/k, n, r, s), n \geq 0\}$ converges to the one for $\{W_k(N/k, n, r), n \geq 0\}$ as $s \rightarrow \infty$. As will be seen in these sections the polynomials

$$\varphi_k(n, x) = \sum_{l=0}^{[n/k]} \binom{n-kl}{l} (-x)^n, \quad n \geq 0 \quad (2)$$

take an important place in our theory.

In Section 4, we are concerned with the generating functions for the particular case $k=1$. For $k=1$, the polynomials $\varphi_1(n, x)$ are closely related to the Chebyshev polynomials of the second kind. C.J. Everett et al. [7] has sought the roots of $\varphi_1(n, x)$, in order to evaluate the number of random walk paths between two absorbing barriers, which is given by $T_1(N, n, 0, 0)$ according to our notation. So as to evaluate the number $T_1(N, n, r, s)$, we express the generating function in terms of Chebyshev polynomials and expand it. As a result, we find an explicit expression for $T_1(N, n, r, s)$ which has a different form from (1).

2. Generating function for $\{W_k(N/k, n, r), n \geq 0\}$

As a preparation for deriving the generating functions for $\{W_k(N/k, n, r), n \geq 0\}$, let us quote the following two lemmas:

Lemma 1 (Pólya and Szegő [8], Gould [9]). *If a function $u(x)$ satisfies the equation $u = 1 + xu^\beta$ for an integer β and $u(0) = 1$, then the following expansions are valid for any integer α :*

$$u^\alpha = \sum_{n=0}^{\infty} A_n(\alpha, \beta) x^n, \quad \alpha \neq 0 \quad (3)$$

$$\frac{u^{\alpha+1}}{(1-\beta)u + \beta} = \sum_{n=0}^{\infty} \binom{\alpha + \beta n}{n} x^n \quad (4)$$

for $|x| < |(\beta - 1)^{\beta-1}/\beta^\beta|$, where

$$A_n(\alpha, \beta) = \frac{\alpha}{\alpha + \beta n} \binom{\alpha + \beta n}{n}, \quad n \geq 0 \quad (5)$$

Lemma 2 (Gould [9]). *For integers α_1, α_2 and β , the following convolution of Vandermonde's type is valid:*

$$\sum_{l=0}^n \binom{\alpha_1 + \beta l}{l} A_{n-l}(\alpha_2, \beta) = \binom{\alpha_1 + \alpha_2 + \beta n}{n} \quad (6)$$

where $A_m(\alpha_2, \beta)$ is defined by (5) for $m = 0, \dots, n$.

In our previous paper [6], we have expressed $W_k(N/k, n, r)$ in terms of $\{A_l(r+1, k+1)\}$ defined by (5) with $\alpha = r+1$ and $\beta = k+1$ as follows:

Lemma 3 (T. Cong and Sato [6]). *For $N, n, r \geq 0$ and $k \geq 1$, $W_k(N/k, n, r)$ is given by*

$$W_k\left(\frac{N}{k}, n, r\right) = \begin{cases} \binom{(k+1)n - N + r}{n}, & 0 \leq n < \left\lfloor \frac{N}{k} \right\rfloor, \\ \sum_{l=0}^{\lfloor N/k \rfloor} \binom{(k+1)l - N - 1}{l} A_{n-l}(r+1, k+1), & n \geq \left\lfloor \frac{N}{k} \right\rfloor + 1. \end{cases} \quad (7)$$

$$(8)$$

Note that if $N \geq r+1$, we define, for convenience, $W_k(N/k, n, r)$ by the right hand side of (7) for $0 \leq n \leq [(N-r-1)/k]$.

Defining the generating function

$$W_k\left(\frac{N}{k}, x, r\right) = \sum_{n=0}^{\infty} W_k\left(\frac{N}{k}, n, r\right) x^n, \quad N, r \geq 0,$$

then we have the following:

Theorem 1. For $N, r \geq 0$, the generating functions of $W_k(N/k, n, r)$, $n \geq 0$, are given by

$$W_k(N/k, x, r) = u^{r+1} \varphi_k(N, x) \quad (9)$$

where $\varphi_k(N, x)$ is defined by (2) and the function u is defined in Lemma 1 with $\beta = k+1$.

Proof. For $n=0, \dots, [N/k]$, the right hand side of (8) becomes, using the convolution (6),

$$\begin{aligned} \sum_{l=0}^{[N/k]} \binom{(k+1)l-N-1}{l} A_{n-l}(r+1, k+1) &= \sum_{l=0}^n \binom{(k+1)l-N-1}{l} A_{n-l}(r+1, k+1) \\ &= \binom{(k+1)n-N+r}{n} \end{aligned}$$

where $A_n(r+1, k+1) = 0$ for $n < 0$. Therefore equation (8) is valid for $n \geq 0$, so that we have

$$\begin{aligned} W_k\left(\frac{N}{k}, x, r\right) &= \sum_{n=0}^{\infty} \sum_{l=0}^{[N/k]} \binom{(k+1)l-N-1}{l} A_{n-l}(r+1, k+1) x^n \\ &= \sum_{l=0}^{[N/k]} \binom{(k+1)l-N-1}{l} \sum_{n=l}^{\infty} A_{n-l}(r+1, k+1) x^n \\ &= \sum_{l=0}^{[N/k]} \binom{(k+1)l-N-1}{l} x^l \sum_{n=0}^{\infty} A_n(r+1, k+1) x^n \\ &= \sum_{l=0}^{[N/k]} \binom{N-kl}{l} (-x)^l \sum_{n=0}^{\infty} A_n(r+1, k+1) x^n. \end{aligned}$$

Putting $\alpha = r+1$ and $\beta = k+1$ in the expansion (3), we find finally

$$W_k(N/k, x, r) = u^{r+1} \varphi_k(N, x),$$

where the function u satisfies the equation $u = 1 + xu^{k+1}$ and $\varphi_k(N, x)$ is defined by (2). \square

Corollary 1. For $N \geq r+1$,

$$\sum_{n=[(N-r-1)/k]+1}^{\infty} W_k\left(\frac{N}{k}, n, r\right) x^n = u^{r+1} \varphi_k(N, x) - \varphi_k(N-r-1, x) \quad (10)$$

where the function u satisfies the equation $u = 1 + xu^{k+1}$ and $\varphi_k(n, x)$ is defined by (2).

Proof. From (7), it follows immediately that

$$\sum_{n=0}^{[(N-r-1)/k]} W_k\left(\frac{N}{k}, n, r\right) x^n = \sum_{n=0}^{[(N-r-1)/k]} \binom{(k+1)n - N + r}{n} x^n = \varphi_k(N-r-1, x),$$

which, in combination with (9), completes the proof. \square

3. Generating function for $\{T_k(N/k, n, r, s), n \geq 0\}$

Now consider a minimal lattice path restricted by two parallel straight lines with an incline k (≥ 1), with which we are particularly concerned in this paper.

$T_k(N/k, n, s+N, s)$ is the number of minimal paths from $(0, 0)$ to $(n, kn+s)$ without crossing any of the lines $y = k(x - N/k)$ and $y = kx + s$ for $N, n, s \geq 0$ (see Fig. 3). Let

$$T_k\left(\frac{N}{k}, x, s+N, s\right) = \sum_{n=0}^{\infty} T_k\left(\frac{N}{k}, n, s+N, s\right) x^n$$

be the generating function for $\{T_k(N/k, n, s+N, s), n \geq 0\}$.

Theorem 2. For $N, s \geq 0$,

$$T_k\left(\frac{N}{k}, x, s+N, s\right) = \frac{\varphi_k(N, x)}{\varphi_k(N+s+1, x)} \quad (11)$$


where $\varphi_k(n, x)$ is defined by (2).

Proof. For $n = 0, \dots, [N/k]$, $T_k(N/k, n, s+N, s)$ is the number of minimal lattice paths from $(0, 0)$ to $(n, kn+s)$ without crossing the line $y = kx + s$. This is given by $W_k(0, n, s)$, that is,

$$T_k(N/k, n, s+N, s) = A_n(s+1, k+1) \quad \text{for } 0 \leq n \leq [N/k]. \quad (12)$$

While, for $n \geq [N/k] + 1$, let us consider a minimal lattice path from $(0, 0)$ to $(n, kn+s)$ crossing the line $y = kx + s$ but not the line $y = k(x - N/k)$. $T_k(N/k, l, s+N, s)$ is the number of minimal lattice paths from $(0, 0)$ to $(l, kl+s)$ without crossing any of the lines $y = k(x - N/k)$ and $y = kx + s$, and $W_k((s+N+1)/k, n-l, s+N)$ is the number of minimal lattice paths from $(l, kl+s+1)$ to $(n, kn+s)$ without crossing the line $y = k(x - N/k)$ for $l = 0, 1, \dots, n-1$. Therefore

$$\sum_{l=0}^{n-1} T_k\left(\frac{N}{k}, l, s+N, s\right) W_k\left(\frac{s+N+1}{k}, n-l, s+N\right)$$

Fig. 3. $T_k(N/k, n, s+N, s)$.

is the number of minimal lattice paths from $(0, 0)$ to $(n, kn+s)$ crossing the line $y=kx+s$ but not crossing the line $y=k(x-N/k)$. Hence we have the following recurrence relation for $T_k(N/k, n, s+N, s)$:

$$T_k\left(\frac{N}{k}, n, s+N, s\right) = W_k\left(\frac{N}{k}, n, s+N\right) - \sum_{l=0}^{n-1} T_k\left(\frac{N}{k}, l, s+N, s\right) W_k\left(\frac{s+N+1}{k}, n-l, s+N\right)$$

for $n \geq [N/k] + 1$. Since $W_k((s+N+1)/k, 0, s+N) = 1$ holds for $s, N \geq 0$, we have

$$\sum_{l=0}^n T_k\left(\frac{N}{k}, l, s+N, s\right) W_k\left(\frac{s+N+1}{k}, n-l, s+N\right) = W_k\left(\frac{N}{k}, n, s+N\right) \quad (13)$$

for $n \geq [N/k] + 1$. From (6), (7) and (12), the above equation (13) is valid for $n \geq 0$, which leads to

$$\begin{aligned} \left(\sum_{n=0}^{\infty} T_k \left(\frac{N}{k}, n, s+N, s \right) x^n \right) \left(\sum_{n=0}^{\infty} W_k \left(\frac{s+N+1}{k}, n, s+N \right) x^n \right) \\ = \sum_{n=0}^{\infty} W_k \left(\frac{N}{k}, n, s+N \right) x^n. \end{aligned}$$

By appealing to Theorem 1, we have

$$T_k \left(\frac{N}{k}, x, s+N, s \right) = \frac{u^{s+N+1} \varphi_k(N, x)}{u^{s+N+1} \varphi_k(s+N+1, x)} = \frac{\varphi_k(N, x)}{\varphi_k(s+N+1, x)}. \quad \square$$

Let us define the generating functions

$$T_k \left(\frac{N}{k}, x, r, s \right) = \sum_{n=0}^{\infty} T_k \left(\frac{N}{k}, n, r, s \right) x^n$$

for $N, s \geq 0$ and $0 \leq r \leq s+N$. Note that if $N \geq r+1$, we define, for convenience,

$$T_k(N/k, n, r, s) = W_k(N/k, n, r) \quad \text{for } 0 \leq n \leq [(N-r-1)/k]. \quad (14)$$

Theorem 3. For $N, s \geq 0$ and $0 \leq r \leq s+N$, we have

$$T_k \left(\frac{N}{k}, x, r, s \right) = \frac{\varphi_k(N, x) \varphi_k(s+N-r, x)}{\varphi_k(s+N+1, x)} \quad (15)$$

where $\varphi_k(n, x)$ is defined by (2).

Proof. For $n = \max(0, [(N-r-1)/k] + 1), \dots, [(s+N-r)/k] - 1, [(s+N-r)/k]$, $T_k(N/k, n, r, s)$ is the number of minimal lattice paths from $(0, 0)$ to $(n, kn - N + r)$ without crossing the line $y = k(x - N/k)$, which is given by $W_k(N/k, n, r)$. If $N \geq r+1$, (14) is valid for $0 \leq n \leq [(N-r-1)/k]$ as mentioned above. Hence, we have

$$T_k(N/k, n, r, s) = W_k(N/k, n, r), \quad 0 \leq n \leq n_0 - 1, \quad (16)$$

where $[(s+N-r)/k] = n_0 - 1$.

While, for $n \geq n_0$, let us consider a minimal lattice path from $(0, 0)$ to $(n, kn - N + r)$ crossing the line $y = kx + s$ but not the line $y = k(x - N/k)$. $T_k(N/k, l, s+N, s)$ is the number of minimal lattice paths from $(0, 0)$ to $(l, kl + s)$ without crossing any of the lines $y = k(x - N/k)$ and $y = kx + s$, and $W_k((s+N+1)/k, n-l, r)$ is the number of minimal lattice paths from $(l, kl + s + 1)$ to $(n, kn - N + r)$ without crossing the line $y = k(x - N/k)$ for $l = 0, \dots, n - n_0$.

Consequently, we have

$$T_k\left(\frac{N}{k}, n, r, s\right) = W_k\left(\frac{N}{k}, n, r\right) - \sum_{l=0}^{n-n_0} T_k\left(\frac{N}{k}, l, s+N, s\right) W_k\left(\frac{s+N+1}{k}, n-l, r\right) \quad (17)$$

for $n \geq n_0$. From (16) and (17), it follows that

$$\begin{aligned} T_k\left(\frac{N}{k}, x, r, s\right) &= \sum_{n=0}^{\infty} W_k\left(\frac{N}{k}, n, r\right) x^n \\ &\quad - \sum_{n=n_0}^{\infty} \sum_{l=0}^{n-n_0} T_k\left(\frac{N}{k}, l, s+N, s\right) W_k\left(\frac{s+N+1}{k}, n-l, r\right) x^n \\ &= \sum_{n=0}^{\infty} W_k\left(\frac{N}{k}, n, r\right) x^n \\ &\quad - \sum_{l=0}^{\infty} T_k\left(\frac{N}{k}, l, s+N, s\right) \sum_{n=n_0+l}^{\infty} W_k\left(\frac{s+N+1}{k}, n-l, r\right) x^n \\ &= \sum_{n=0}^{\infty} W_k\left(\frac{N}{k}, n, r\right) x^n - \left(\sum_{l=0}^{\infty} T_k\left(\frac{N}{k}, l, s+N, s\right) x^l \right) \\ &\quad \times \left(\sum_{n=n_0}^{\infty} W_k\left(\frac{s+N+1}{k}, n, r\right) x^n \right). \end{aligned}$$

Applying Theorems 1, 2 and Corollary 1, we finally reach

$$\begin{aligned} T_k\left(\frac{N}{k}, x, r, s\right) &= u^{r+1} \varphi_k(N, x) \\ &\quad - \frac{\varphi_k(N, x)}{\varphi_k(s+N+1, x)} \{u^{r+1} \varphi_k(s+N+1, x) - \varphi_k(N+s-r, x)\} \\ &= \frac{\varphi_k(N, x) \varphi_k(s+N-r, x)}{\varphi_k(s+N+1, x)} \end{aligned}$$

where the function u is defined by Theorem 1. \square

As the right hand side of (15) is a ratio of two polynomials in x , explicit expression for $T_k(N/k, n, r, s)$ can be obtained by expanding it. For instance, $\varphi_k(n, x)^{-1}$ can be expanded in the following form:

$$\frac{1}{\varphi_k(n, x)} = \sum_{m=0}^{\infty} \left[\sum_{\substack{l_1 + \dots + l_j = m \\ 1 \leq l_i \leq \lfloor n/k \rfloor, i=1, \dots, j}} (-1)^{m+j} \binom{n-kl_1}{l_1} \dots \binom{n-kl_j}{l_j} \right] x^m \quad (18)$$

Since the generating function (15) for $N=0$ and $r=s$ reduces to

$$T_k(0, x, s, s) = \frac{1}{\varphi_k(s+1, x)},$$

and together with (18), the explicit expression of $T_k(0, n, s, s)$ is given by

$$T_k(0, n, s, s) = \sum_{\substack{l_1 + \dots + l_j = n \\ 1 \leq l_i \leq [(s+1)/k], i=1, \dots, j}} (-1)^{n+j} \binom{s+1-kl_1}{l_1} \dots \binom{s+1-kl_j}{l_j}.$$

Now in order to show that the generating function $T_k(N/k, x, r, s)$ converges to $W_k(N/k, x, r)$ as $s \rightarrow \infty$, we notice:

Lemma 3. For $k \geq 1$ and $|x| < k^k/(k+1)^{k+1}$

$$\lim_{n \rightarrow \infty} \varphi_k(n, x) u^{n+1} = \frac{u}{(k+1) - ku}, \quad (19)$$

where the function u is defined by Theorem 1.

Proof. Let a function u satisfy the equation $u = 1 + xu^\beta$ with $\beta = k+1$ and $u(0) = 1$. From Lemma 1, we have

$$u^{n+1} = \sum_{l=0}^{\infty} A_l(n+1, \beta) x^l, \quad \frac{u^{-n}}{(1-\beta)u + \beta} = \sum_{l=0}^{\infty} \binom{\beta l - n - 1}{l} x^l$$

for $n \geq 0$. From these relations, it follows that

$$\begin{aligned} & \frac{u}{(1-\beta)u + \beta} \\ &= \frac{u^{-n}}{(1-\beta)u + \beta} u^{n+1} = \sum_{l=0}^{[n/k]} \binom{\beta l - n - 1}{l} x^l u^{n+1} + \sum_{l=[n/k]+1}^{\infty} \binom{\beta l - n - 1}{l} x^l u^{n+1} \\ &= \sum_{l=0}^{[n/k]} \binom{\beta l - n - 1}{l} x^l u^{n+1} + \sum_{l=[n/k]+1}^{\infty} \left(\sum_{m=[n/k]+1}^l \binom{\beta m - n - 1}{m} A_{l-m}(n+1, \beta) \right) x^l. \end{aligned} \quad (20)$$

Applying the convolution (6), the inequality

$$\begin{aligned} & \sum_{m=[n/k]+1}^l \binom{\beta m - n - 1}{m} A_{l-m}(n+1, \beta) \\ & < \sum_{m=0}^l \binom{\beta m - n - 1}{m} A_{l-m}(n+1, \beta) = \binom{\beta l}{l}, \quad n \geq 0 \end{aligned}$$

follows for $l \geq [n/k] + 1$. Since

$$\lim_{n \rightarrow \infty} \sum_{l=[n/k]+1}^{\infty} \binom{\beta l}{l} x^l = 0$$

holds for $|x| < |(\beta-1)^{\beta-1}/\beta^\beta|$, the second term of the right hand side of (20) converges to 0. Therefore we have

$$\frac{u}{(1-\beta)u + \beta} = \lim_{n \rightarrow \infty} \left[\sum_{l=0}^{[n/k]} \binom{\beta l - n - 1}{l} x^l \right] u^{n+1}.$$

Putting $\beta = k + 1$, the above equation gives (19). \square

After this preparation, we have

Theorem 4. For $N, r \geq 0$ and $k \geq 1$,

$$\lim_{s \rightarrow \infty} T_k\left(\frac{N}{k}, x, r, s\right) = W_k\left(\frac{N}{k}, x, r\right), \quad |x| < \frac{k^k}{(k+1)^{k+1}}.$$

Proof. From Theorems 1, 2 and Lemma 3, we have

$$\begin{aligned} \lim_{s \rightarrow \infty} T_k\left(\frac{N}{k}, x, r, s\right) &= \lim_{s \rightarrow \infty} \frac{\varphi_k(N, x) \varphi_k(s + N - r, x)}{\varphi_k(s + N + 1, x)} \\ &= \lim_{s \rightarrow \infty} \frac{u^{s+N-r+1} \varphi_k(s + N - r, x)}{u^{s+N+2} \varphi_k(s + N + 1, x)} \varphi_k(N, x) u^{r+1} \\ &= u^{r+1} \varphi_k(N, x) = W_k\left(\frac{N}{k}, x, r\right). \quad \square \end{aligned}$$

Next, we show that there is a difference equation which the polynomial $\varphi_k(n, x)$ satisfies, in the following:

Theorem 5. The polynomial $\varphi_k(n, x)$ satisfies the difference equation

$$\varphi_k(n, x) = \begin{cases} \varphi_k(n-1, x) - x\varphi_k(n-k-1, x), & n \geq k+1, \\ 1, & 0 \leq n \leq k. \end{cases} \quad (21)$$

Proof. This is clear since

$$\sum_{n=0}^{\infty} \varphi_k(n, x) z^n = 1/(1 - z + xz^{k+1}). \quad \square$$

4. Explicit expression for $T_1(N, n, r, s)$

In this section, we consider the particular case $k = 1$. In this case, $W_1(N, n, r)$ is the number of minimal lattice paths from $(0, 0)$ to $(n, n - N + r)$ without crossing the line $y = x - N$ for $n, N, r \geq 0$ and $n \geq N - r$, which is simply given by

$$W_1(N, n, r) = \binom{2n - N + r}{n} - \binom{2n - N + r}{n - N - 1}, \quad n \geq N - r$$

by means of the reflection principle. $T_1(N, n, r, s)$ is the number of minimal lattice paths from $(0, 0)$ to $(n, n - N + r)$ without crossing any of the lines $y = x - N$ and $y = x + s$ for $n, N, s \geq 0$, $0 \leq r \leq s + N$ and $n \geq N - r$. In order to evaluate $T_1(N, n, r, s)$, we express the generating functions in terms of Chebyshev polynomials.

Theorem 6. (i) For $N, r \geq 0$,

$$W_1(N, x, r) = \left(\frac{1 - \sqrt{1 - 4x}}{2x} \right)^{r+1} x^{N/2} U_N \left(\frac{1}{2\sqrt{x}} \right).$$

(ii) For $N, s \geq 0$ and $0 \leq r \leq s + N$,

$$T_1(N, x, r, s) = x^{(N-r-1)/2} \frac{U_N(1/2\sqrt{x}) U_{s+N-r}(1/2\sqrt{x})}{U_{s+N+1}(1/2\sqrt{x})} \quad (22)$$

where $U_n(x)$ is Chebyshev polynomial of the second kind

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta}, \quad x = \cos \theta.$$

Proof. (i) and (ii) are trivial since the polynomial $\varphi_1(n, x)$ can be written as:

$$\varphi_1(n, x) = \sum_{l=0}^{[n/2]} \binom{n-l}{l} (-x)^l = x^{n/2} U_n \left(\frac{1}{2\sqrt{x}} \right), \quad n \geq 0. \quad \square$$

In virtue of Theorem 6(ii), explicit expression for $T_1(N, n, r, s)$ can be obtained as follows:

Theorem 7. For $N, n, s \geq 0$ and $0 \leq r \leq s + N$,

$$T_1(N, n, r, s) = \frac{4}{s + N + 2} \sum_{\nu=1}^M (2 \cos \theta_\nu)^{2N+N} \sin(N+1)\theta_\nu \sin(r+1)\theta_\nu \quad (23)$$

where $\theta_\nu = \nu\pi/(s + N + 2)$, $\nu = 1, \dots, [\frac{1}{2}(s + N + 1)] (= M)$.

Proof. Let us put

$$x = (2 \cos \theta)^{-2}, \quad 0 < \theta < \frac{1}{2}\pi.$$

From (22), it follows that

$$T_1(N, x, r, s) = (2 \cos \theta)^{-(N-r-1)} \frac{\sin(N+1)\theta \sin(s+N-r+1)\theta}{\sin \theta \sin(s+N+2)\theta}.$$

The θ are $[\frac{1}{2}(s + N + 1)]$ different roots of $\sin(s + N + 2)\theta$:

$$\theta_\nu = \frac{\nu\pi}{s + N + 2}, \quad \nu = 1, \dots, [\frac{1}{2}(s + N + 1)] (= M)$$

which imply M different roots of $\varphi_1(s + N + 1, x)$ such that

$$x_\nu = (2 \cos \theta_\nu)^{-2}, \quad \nu = 1, \dots, M.$$

Consequently, the coefficient of x^n in the expansion of $T_1(N, x, r, s)$ is represented

by

$$\sum_{\nu=1}^M (2 \cos \theta_{\nu})^{2n-N+r+3} \frac{\sin(N+1)\theta_{\nu} \sin(s+N-r+1)\theta_{\nu}}{(-\sin \theta_{\nu})(s+N+2)(\cos(s+N+2)\theta_{\nu})[d\theta/dx]_{\theta=\theta_{\nu}}}$$

$$= \frac{4}{s+N+2} \sum_{\nu=1}^M (2 \cos \theta_{\nu})^{2n+r-N} \sin(N+1)\theta_{\nu} \sin(r+1)\theta_{\nu},$$

which give $T_1(N, n, r, s)$. \square

The finite sum (1) as mentioned in Section 1 is equal to $T_1(a-1, n, a-1-i, b-1)$, where $n = \frac{1}{2}(t+i)$. From (23), it is immediately shown that

Corollary 2. For $t+i \equiv 0 \pmod{2}$ and $-b < i < a$,

$$\sum_{l=-\infty}^{\infty} \left\{ \binom{t}{\frac{1}{2}(t+i)+l(a+b)} - \binom{t}{\frac{1}{2}(t+i)+b+l(a+b)} \right\}$$

$$= \frac{4}{a+b} \sum_{\nu=1}^M (2 \cos \theta_{\nu})^t \sin a\theta_{\nu} \sin(a-i)\theta_{\nu},$$

where $\theta_{\nu} = \nu\pi/(a+b)$, $\nu = 1, 2, \dots, [\frac{1}{2}(a+b-1)] (=M)$.

Acknowledgement

The authors wish to express their gratitude to Professor T. Kitagawa, the director of the International Institute for Advanced Study of Social Information Science, Fujitsu, for his suggestions. We are also grateful to Professor T. Ishihara for his encouragement and helpful comments.

References

- [1] S. Dua, S. Khadilkar and K. Sen, A modified random walk in the presence of partially reflecting barriers, *J. Appl. Prob.* 13 (1976) 169–175.
- [2] D.G. Rogers, A Schröder triangle: three combinatorial problems in: *Combinatorial Mathematics V*, Proc. Fifth Austrian Conf., Lecture Notes in Mathematics 622 (Springer-Verlag, Berlin, 1977) 175–196.
- [3] D.G. Rogers, Pascal triangles, Catalan numbers and renewal arrays, *Discrete Math.* 22 (1978) 301–310.
- [4] D.A. Klarner, Correspondence between plane trees and binary sequences, *J. Combin. Theory* 9 (1970) 401–411.
- [5] A.D. Sands, On generalized Catalan numbers, *Discrete Math.* 21 (1978) 219–221.
- [6] T.T. Cong and M. Sato, One-dimensional random walk with unequal step lengths restricted by an absorbing barrier, *Discrete Math.* 40 (1982) 153–162.
- [7] C.J. Everett and P.R. Stein, The combinatorics of random walk with absorbing barriers, *Discrete Math.* 17 (1977) 27–45.
- [8] G. Polya and G. Szegő, *Aufgaben und Lehrsätze aus der Analysis*, Vol. 1 (Springer-Verlag, Berlin, 1925).
- [9] H.W. Gould, Some generalizations of Vandermonde's convolution, *Amer. Math. Monthly* 63 (1956) 84–91.